NOTES ON NEW AND NOTEWORTHY PLANTS. LXXX

Harold N. Moldenke

CASSELIA GLAZIOVII var. SERRATA Moldenke, var. nov.

Hasc varietas a forma typica speciei laminis foliorum uniforme

regularitorque serratis differt.

This variety differs from the typical form of the species in having the margins of its leaf-blades merely uniformly and regularly serrate with comparatively small acute or subacute antrorse teeth.

The type of the variety was collected by G. Hatschbach, W. Anderson, R. Barneby, and B. Gates (no. 36255) on a "campo limpo" 12 km. north of São João da Alianca, Goiás, Brazil, on February 18, 1975, and is deposited in my personal herbarium at Plainfield, New Jersey. The corollas are described by the collectors as having been lilac in color.

MNEIFFIA FRUTICOSA var. UNGUICULATA (Fern.) Moldenke, comb. nov.

Oenothera fruticosa var. unguiculata Fern., Rhodora 11: 551,
pl. 577, fig. 1-3. 1939.

LANTANA BAHAMENSIS var. FLORIDANA Moldenke, var. nov.

Haec varietas a forma typica speciei petiolis plerumque pro-

portione brevioribus recedit.

This variety differs from the typical form of the species in having the petioles on mature leaves usually proportionately shorter.

The type of the variety was collected by John Kunkel Small ($\underline{\text{no. 2101}}$) on the beach opposite Miami, Dade County, Florida, in November, 1904, and is deposited in the Britton Herbarium at the New York Botanical Garden.

STACHYTARPHETA CONFERTIFOLIA var. PUBERULENTA Moldenke, var. nov. Haec varietas a forma typica speciei ramis ubique densiuscule puberulis differt.

This variety differs from the typical form of the species in having its stems and branches rather densely puberulent throughout.

The type of the variety was collected by G. Hatschbach, W. Anderson, R. Barneby, and B. Gates (no. 36361) on a sandy campo, at an altitude of 1300 to 1500 meters, GO.12, Chapada da dos Veadeiros, in the municipality of Alto Paraiso, Goiás, Brazil, on February 20, 1975, and is deposited in my personal herbarium at Plainfield, New Jersey. The corollas are described by the collectors as having been white.